

Anesthetic Management of a Parturient with a Syringo-Pleural Shunt

Shuaib K., Koh J., Factor M., Angelo T.E., Daoud B.E.

Introduction:

- Impaired CSF drainage in Arnold Chiari Malformation (ACM) leads to high cranial pressures being transmitted to the central canal.¹
- The development of syringomyelia is characterized by a fluid filled cavity in the spinal cord which is frequently associated with ACM.²
- Shunt placement through a syrinx is indicated for patients who failed decompression due to syrinx enlargement or persistent neurologic symptoms.³

Case:

- 29yo G3P0 at 31wks gestational age with a h/o ACM type II w/ associated syringomyelia now s/p neurosurgery & syringo-pleural shunt placement.
- Patient admitted for a mandibular abscess with an associated 12kg weight gain during her stay. CXR revealed bilateral pleural opacities. Repeat episodes of fetal bradycardia with minimal variability.
- Required emergent cesarean delivery general anesthesia.
- Persistently hypoxic (SpO2 87%) post-operatively. A CT-angiogram of chest revealed bilateral multifocal pneumonia. Shuntogram revealed no kinking or obstruction of her syringo-pleural shunts




Figure 1: Spine magnetic resonance imaging of a parturient with syringomyelia. Cervicothoracic syrinx as well as associated seroma at C6-C7 level seen on T2 weighted images.

Question 1: Which anesthetic technique is more favorable ?

General Anesthesia⁴:

- Better management of ↑ ICP through BP control and hyperventilation
- Avoids risk of dural puncture

VS

Neuraxial Anesthesia⁵:

- Avoids ↑ ICP from laryngeal manipulation and intubation
- Avoids narcotic effects on fetus

Question 2: What mode of delivery is more favorable ?

Cesarean Delivery with GA⁴:

- Avoids changes in ICP that occur with Valsalva & uterine contractions
- Avoids risk of dural puncture

VS

Vaginal Delivery⁵:

- Can slowly titrate labor epidural to prevent acute reductions in BP
- Avoids manipulation of airway and augmentation of pulmonary function.

OBSTETRIC ANESTHESIOLOGY: ORIGINAL CLINICAL RESEARCH REPORT

Anesthetic and Obstetric Management of Syringomyelia During Labor and Delivery: A Case Series and Systematic Review


Garvey, Gráinne Patricia MBChB (euro), FRCA^{*}; Wasade, Vibhangini S. MD[†]; Murphy, Kellie E. MD^{‡§}; Balki, Mrinalini MD^{*‡}

[Author Information](#)👍

Anesthesia & Analgesia: September 2017 - Volume 125 - Issue 3 - p 913-924



Renaissance School of Medicine
Stony Brook University



Results⁶:

- Prior to delivery: worsening neurological condition in 5 patients (11.6%), 4 of whom underwent GA.
- Peripartum: one neurological complication (one sided anesthesia after delivery)
- No persistent complications from spontaneous vaginal delivery & neuraxial anesthesia.

Takeaways:

1. Physiologic changes in pregnancy such as increased ICP make syringomyelia a significant clinical concern in pregnant patients.
2. Unique problems such as neuro-shunt malfunction that can occur after many years of excellent function add another dimension of complexity in those with syringo-pleural shunts⁷
3. A shunt patency study is required if there is suspicion of a malfunctioning shunt
4. It appears that anesthetic complications occur infrequently regardless of anesthetic technique
5. It appears that neuraxial anesthesia might be undertaken without serious clinical consequence in these patients

References:

1. Williams B. Further Thoughts on the Valvular Action of the Arnold-Chiari Malformation. *Developmental Medicine & Child Neurology*, 1971 Volume 13: 105-112
2. Garvey GP, Wasade VS, Murphy KE, Balki M. Anesthetic and Obstetric Management of Syringomyelia During Labor and Delivery: A Case Series and Systematic Review, *Anesthesia & Analgesia*: September 2017 - Volume 125 - Issue 3 - p 913-924
3. Schijman E, Steinbok P. International survey on the management of Chiari I malformation and syringomyelia. *Childs Nerv Syst*. 2004 May;20(5):341-8. Epub 2004 Feb 14. PMID: 14966661.
4. Ghaly RF, Candido KD, Sauer R, Knezevic NN. Anesthetic management during Cesarean section in a woman with residual Arnold-Chiari malformation Type I, cervical kyphosis, and syringomyelia. *Surg Neurol Int* 2012;3:26
5. Roelofse JA, Shipton EA, NellAC. Anesthesia for caesarean section in a patient with syringomyelia. *SA Med J* 1984; 65: 736-7.
6. Garvey GP, Wasade VS, Murphy KE, Balki M. Anesthetic and Obstetric Management of Syringomyelia During Labor and Delivery: A Case Series and Systematic Review. *Anesthesia and analgesia*. 2017;125(3)
7. Wisoff JH, Kratzert KJ, Handwerker SM, Young BK, Epstein F. Pregnancy in Patients with Cerebrospinal Fluid Shunts: Report of a Series and Review of the Literature, *Neurosurgery*, 1991;29: 827-831