

Development and Validation of a Prognostic Model for Postpartum Haemorrhage

B. O'Shea, B. McCaffrey, J. Lloyd, N. Nguyen-Lu and D. J. N. Wong

Background

- Difficult to predict and no widely accepted prognostic tools
- Perform poorly or specific sub-groups/specific biomarkers

Prognostic Tool Development

- Risk factors derived from literature search
- Odds Ratios for risk factors converted to regression coefficient weights (Moon et al)
- Weights adjusted by author consensus
- Scores ascribed to each risk factor

Uncomplicated pregnancy	0
Singleton	0
Elective C-section	1
Induction/augmentation of labour	1
Lower segment fibroids	1
Multiple pregnancy	1
Previous myomectomy	1
Emergency caesarean-section or caesarean in labour	2
Platelets $<100 \times 10^9/L$	2
Instrumental birth	2
Known coagulopathy	2
Previous caesarean section or uterine surgery	2
Chorioamnionitis/sepsis in labour	2
Macrosomia $>4000g$	2
Placenta previa (posterior)	2
Placenta accreta	3

Methods and Results

Retrospective cohort of women giving birth over 1 year (2021) in tertiary centre

Included: 6334 women, all modes of delivery, emergency/elective


Data collected using electronic health records (Badgernet)

ROC curve analysis to assess discriminative performance

Calibration of score: estimated probabilities of PPH plotted against PPH scores

Performance tested over 3 definitions of PPH: $\geq 500\text{ml}$, $\geq 1000\text{ml}$ and $\geq 1500\text{ml}$ blood loss

Blood Loss	ROC Curve - AUC
$\geq 500\text{ml}$	0.719 (95% CI: 0.707-0.731)
$\geq 1000\text{ml}$	0.661 (95% CI: 0.644- 0.678)
$\geq 1500\text{ml}$	0.661 (95% CI: 0.633-0.689)



Key Points

- Performed well over a variety of PPH definitions
- Simple to use tool and applicable across broad obstetric population
- Aids risk stratification and may guide strategies to reduce blood loss
- Limitations: Single centre, tertiary cohort, low number at highest risk

Next steps

- Assess changes to clinical care following local implementation
- External validation