



Quadratus Lumborum in Cesarean Section Trial (QUALICS)

Hermann dos Santos Fernandes, MD PhD^{1, 2}; Fernando Nani, MD¹; Shirley Andrade Santos, MD PhD^{1, 2}; Rossana Francisco, MD PhD¹; Joaquim Edson Vieira, MD PhD¹;


Cristian Arzola*, MD MSc²

1. Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil

2. Mount Sinai Hospital, University of Toronto, Canada



Anesthesiology & Pain Medicine
UNIVERSITY OF TORONTO



**Mount Sinai
Hospital**

INTRODUCTION

- Postoperative pain after Cesarean Section (CS) has significant impact on quality of recovery
- Neuraxial morphine is the standard for post-CS analgesia with potential adverse effects
- Alternatives to neuraxial morphine
 - ✓ TAP
 - Inferior analgesic effect than neuraxial morphine
 - ✓ **QLB**
 - Superior to TAP
 - ✓ Is QLB equivalent to neuraxial morphine?

Objective: to evaluate the post-CS analgesic efficacy of QLB, in comparison with both intrathecal morphine (ITM) and ITM associated with QLB

METHODS


- A randomized blinded clinical trial
- 57 term, healthy parturients scheduled to elective CS.
- Three groups:
 - ITM group :::: spinal morphine with no QLB
 - QLB group :::: QLB with no ITM
 - QLB-ITM group :::: both QLB and ITM
- Standard spinal anesthesia:
 - ✓ 0.5% hyperbaric bupivacaine - 12.5 mg
 - ✓ Fentanyl - 20 mcg
 - ✓ Morphine - 80 mcg (ITM and QLB-ITM groups)
 - ✓ Placebo (0.4 ml of 0.9% NaCl) (QLB group)

METHODS

- Primary outcomes:
 - ✓ Pain scores (0-100 at rest and movement) on POD-1
 - ✓ Opioid consumption on POD-1
- Secondary outcomes:
 - ✓ Nausea/vomiting, pruritus, urinary retention and persistent postoperative pain at 3 months

QLB – POSTERIOR APPROACH

- After skin closure and still under residual sensory block
 - Ultrasound-guided bilateral posterior QLB
 - Ropivacaine 0.375%, 20 ml in each side
 - QLB and QLB-ITM groups
 - ITM group:
 - Mock ultrasound scanning performed under the surgical drapes



Recruitment N=61	ITM	QLB	QLB-ITM
Inclusion / Exclusion	17 / 0	21 / 2	23 / 2
Assessment POD-1	17	19	21
Assessment 3- month	13	8	15

RESULTS

CONCLUSIONS

- QLB demonstrated no analgesic difference to ITM, and no added benefit when both are associated (QLB-ITM)
- QLB provided the benefit of causing less urinary retention
- QLB can be an alternative when ITM is not administered without compromising postoperative analgesia

REFERENCES

Roofthoof. PROSPECT guideline for elective caesarean section. Anaesthesia 2021;76:665-80

OUTCOMES	ITM	QLB	QLB-ITM	p-value
Total IV morphine consumption on POD-1 (mg)	3.41 (4.84)	3.10 (2.55)	1.76 (2.96)	0.303
Pain at rest on POD-1 (NVS 0-100)	13.52 (23.43)	18.94 (17.28)	5.71 (14.68)	0.084
Pain at movement on POD-1 (NVS 0-100)	35.29 (30.84)	53.15 (26.04)	41.42 (23.51)	0.131
Urinary retention on POD-1 (%)	7.02	0	1.75	0.032

ANOVA (Bonferroni test), Chi-square and Fisher’s exact test. Values are mean (SD) and percentage (%)