

## Background

- The incidence of obesity in pregnancy is increasing in North America. *(Driscoll A.K., NCHS Data Brief 2020)*
- In patients with obesity, the post-operative period is characterized by an increased and prolonged inflammatory response, making it an independent risk factor for acute pain. *(Mei W, Eur j Pain. 2010)*
- Providers tend to prescribe less opioids to these patients, which may inappropriately undertreat their pain. *(Bui AH, J Pharm Policy Pract. 2018)*
- While obesity is an independent risk factor for acute pain in the non-obstetric surgical population, the impact of obesity on overall quality of postpartum recovery remains under explored.

## Methods


- Single-center prospective cohort study
- Two groups:
  - Control (BMI<30 kg/m<sup>2</sup>) & High BMI (BMI>40 kg/m<sup>2</sup>)
- Standardized intraoperative and postoperative care
- Primary Outcome:
  - Difference in the ObsQoR-10 scores at 24h post-CD
- Secondary outcomes:
  - Difference in pain scores, total opioid consumption, incidence of adverse events, time to first mobilization, length of hospital stay, breastfeeding at 24h and readmission rates

## Hypothesis

We hypothesized that parturients with Class 3 obesity will demonstrate worse quality of inpatient postpartum recovery following elective cesarean delivery as measured by the ObsQoR-10 when compared to non-obese parturients.

Results

|                                 | BMI < 30    | BMI > 40    | Difference (95% CI) | P    |
|---------------------------------|-------------|-------------|---------------------|------|
| Mean (SD) QOR score at 24 hours | 80.1 (12.1) | 80.3 (13.1) | 0.3 (-4.1 to 4.7)   | 0.90 |



| Variable                                        | BMI < 30           | BMI > 40           | RR or difference (95% CI) | P     |
|-------------------------------------------------|--------------------|--------------------|---------------------------|-------|
| Median [IQR] total inpatient opioid consumption | 15 [0 – 30]        | 7.5 [0 – 35]       | -7.5 (-18 to 12)          | 0.78  |
| Time to first opioid request (h)                | 4.4 [3 – 11.5]     | 5 [2 – 15]         | 0.6 (-3 to 8)             | 0.97  |
| Breastfeeding at 24h                            | 56 (90%)           | 42 (70%)           | 0.78 (0.64 to 0.93)       | 0.006 |
| Time to first mobilization(h)                   | 10.3 [8 – 13.8]    | 8 [7.5 – 11.8]     | -2.3 (-4 to -0.5)         | 0.03  |
| Time to 1st intake of non-clear fluids (h)      | 3 [1 – 6.5]        | 2 [1 - 7]          | -1 ( -3 to 1)             | 0.20  |
| Length of stay (h)                              | 49.8 [46.0 – 52.2] | 49.0 [47.3 – 52.8] | -0.8 (-2.1 to 1.7)        | 0.88  |
| Readmission within 6-weeks (%)                  | 5 (8%)             | 2 (4%)             | 0.47 (0.10 to 2.35)       | 0.45  |

# Quality of Recovery in Parturients with Obesity: A Prospective Observational Cohort Study

## Conclusion

- There was **no significant differences in the ObsQOR-10 at 24 hours** between the two groups.
- There was **no significant differences in the NRS pain scores (rest or movement), or opioid consumption** between the two groups.
- Patients with obesity **ambulated sooner** after their surgery, however, were **less likely to breastfeed** at 24 hours.
- Future studies are needed to evaluate quality of outpatient recovery in patients with obesity, including domains such as sleep, fatigue and psychosocial distress.