

Racial Disparities in Maternal and Neonatal Outcomes in the COVID-19 Positive Parturient

Rachel Weeks, MS^{1,2}, Andrea Ruggiero, MS², Louis Ballas², Tungming Leung, PhD², and Gregory Palleschi, MD²

¹Donald and Barbara Zucker School of Medicine at Hofstra/Northwell


²Department of Anesthesia, Long Island Jewish Medical Center

Introduction

- Racial and ethnic disparities exist in maternal healthcare manifesting as higher maternal and neonatal morbidity/mortality among underrepresented minorities.^{2,3}
- Independent of racial and ethnic background, it has been shown by our group that maternal and neonatal outcomes are adversely affected by intrapartum COVID-19 infection.¹
 - A previous study done by our group showed higher rates of ICU admission, ventilator use, pre-eclampsia, HELLP syndrome, preterm labor, cesarean delivery, low APGAR scores, low newborn birthweight, and lower breastfeeding success in the COVID-19 positive parturient and their neonates compared to the COVID-19 negative parturient.
- **Given the known increase in adverse outcomes to those with COVID-19, we sought to study whether underrepresented minorities were disproportionately affected by gestational COVID-19 infection.**

Methods

- **Retrospective chart review**
- Patient population → 2143 parturients who tested positive for COVID-19 during pregnancy and delivered at seven tertiary care hospitals within Northwell Health between March 1, 2020, and June 31, 2021.
- **Divided into 5 racial groups:** White, Black, Asian, Other (Multiracial and Native American), and Unknown
- In this study, we investigated adverse outcomes as defined in previous studies to see if there were differences between these racial groups
 - **Maternal outcomes:** Cesarean section, pre-eclampsia
 - **Neonatal outcomes:** APGAR scores below 7, breastfeeding success
- Chi-squared analyses or Fischer's Exact tests were performed based on sample size eligibility to determine whether the outcomes of interest differed between races in the COVID-19 positive parturient.



Results

- Compared to white women, black women had higher rates of:
 - Cesarean delivery**
 - 38% vs. 28%, $p < 0.001$
 - Pre-eclampsia**
 - 12% vs. 3%, $p < 0.001$
 - APGAR scores less than 7 at 1 minute**
 - 9% vs. 4%, $p < 0.001$
 - APGAR scores less than 7 at 5 minutes**
 - 4% vs. 2%, $p < 0.001$
- Compared to white women, black women had a lower rate of:
 - Breastfeeding success**
 - 21% vs. 28%, $p < 0.001$
- Compared to white women, Asian women had a higher rate of:
 - Cesarean delivery**
 - 37% vs. 28%, $p < 0.001$

Limitations

- Does not control for pre-existing conditions that may have an impact on maternal and neonatal health
 - Includes conditions such as obesity, hypertension, gestational diabetes, etc.
- Prior pre-gestational COVID-19 infection was not included in the analysis
- Did not analyze trimester of infection

Conclusions

- **Underrepresented parturients with gestational COVID-19 and their neonates experience more negative outcomes compared to the majority.**
- Future studies may investigate:
 - ERAS protocol in mitigating disparities between racial and ethnic groups
 - Bias (implicit/explicit)
 - Lack of resources contributing to poor antenatal care
- **Awareness is key!**
 - May help mitigate disparities and provide solutions to prevent negative outcomes in the future

References

1. Katz, J., Ruggiero, A., Ballas, L., Shore-Lesserson, L., & Palleschi, G. (2021). *COVID-19 and the Impact on Pregnancy Outcomes*. Presented at 54th Annual SOAP Meeting.
2. Tangel, V. E., Matthews, K. C., Abramovitz, S. E., & White, R. S. (2020). Racial and ethnic disparities in severe maternal morbidity and anesthetic techniques for obstetric deliveries: A multi-state Analysis, 2007–2014. *Journal of Clinical Anesthesia*, 65, 109821.
3. Tangel, V., White, R. S., Nachamie, A. S., & Pick, J. S. (2018). Racial and ethnic disparities in maternal outcomes and the disadvantage of Peripartum Black Women: A Multistate Analysis, 2007–2014. *American Journal of Perinatology*, 36(08), 835–848.