



Endometriosis of the Lungs Causing Recurrent Pneumothoraces in Pregnancy


Katelyn Scharf MD, Shobana Bharadwaj MBBS, Jessica Galey MD, Autumn LaRocque MD, Bhavani Kodali MBBs; MD
Department of Anesthesiology, University of Maryland School of Medicine, Baltimore, MD

Introduction

- Thoracic endometriosis is the presence of endometrial tissue in the diaphragm, pleura, or lungs
- Though rare, it is the most common form of endometriosis outside of the pelvis
- Affects women of reproductive age
- Thought to be caused by endometrial tissue traveling through diaphragmatic defects or via microembolization in pelvic veins
- Often will manifest as pneumothoraces

Clinical Presentation

37-year-old woman from Ghana with a PMHx of uterine fibroids s/p myomectomy presented for progressive fatigue, shortness of breath, and bilateral pneumothoraces. Initial work-up, including transthoracic needle aspiration of a nodule in her left lower lobe was non-diagnostic. She presented to our institution for a second opinion.





Endometriosis of the Lungs Causing Recurrent Pneumothoraces in Pregnancy

Katelyn Scharf MD, Shobana Bharadwaj MBBS, Jessica Galey MD, Autumn LaRocque MD, Bhavani Kodali MBBs; MD
Department of Anesthesiology, University of Maryland School of Medicine, Baltimore, MD

Imaging



CT chest without contrast revealed a moderate-sized septated right-sided pneumothorax, numerous pulmonary nodules, masses, and cystic lesions distributed throughout all 5-lobes.

Delivery Planning

- Recommended for primary cesarean delivery (CD) given risk of pneumothorax with Valsalva maneuver
- Underwent uncomplicated CD at 36w4d
- During CD, indwelling chest tubes were placed to suction
- CSE was performed without respiratory issues
- Fetus was delivered vertex via Kiwi vacuum assistance to avoid fundal pressure and potential popping of a pulmonary bleb
- 24-hour postpartum monitoring




Endometriosis of the Lungs Causing Recurrent Pneumothoraces in Pregnancy



Katelyn Scharf MD, Shobana Bharadwaj MBBS, Jessica Galey MD, Autumn LaRocque MD, Bhavani Kodali MBBs; MD
Department of Anesthesiology, University of Maryland School of Medicine, Baltimore, MD

Postpartum

- Underwent video-assisted thoracic surgery (VATS) for pleurodesis and wedge biopsy, which confirmed endometriosis of the lungs
- Patient returned to Ghana 2-months postpartum and was lost to follow up



A) Endometriosis of lung, low power: Lung biopsy with endometrial stroma and endometrial glands (center) adjacent to lung parenchyma (periphery), consistent with endometriosis. Inset: high power view of endometrial gland.



B) Endometriosis of lung, high power: Lung biopsy with endometrial stroma and endometrial glands, consistent with endometriosis. Inset: high power view of endometrial gland.

C) Decidualization: lung biopsy with areas of endometrial decidualization, consistent with endometriosis in a pregnant patient. Inset: high power view of decidualized cells.

Discussion

- Multidisciplinary approach that involved MFM, pulmonology, thoracic surgery, and OB anesthesia
- The avoidance of fundal pressure during the C-section helped prevent development of an additional pneumothorax

References

1. Andres MP, Arcoverde FVL, Souza CCC, Fernandes LFC, Abrão MS, Kho RM. Extrapelvic Endometriosis: A Systematic Review. *J Minim Invasive Gynecol.* 2020;27(2):373-389.
2. Rousset P, Rousset-Jablonski C, Alifano M, Mansuet-Lupo A, Buy JN, Revel MP. Thoracic endometriosis syndrome: CT and MRI features. *Clin Radiol.* 2014;69(3):323-330.
3. Channabasavaiah AD, Joseph JV. Thoracic endometriosis: Revisiting the association between clinical presentation and thoracic pathology based on thoracoscopic findings in 110 patients. *Medicine.* 2010;89(3):183-188.