

CESAREAN SECTION WITH EPIDURAL ANESTHESIA IN A MORBIDLY OBESE PARTURIENT WITH POST-COVID PULMONARY FIBROSIS

Yuta Kashiwagi, MD¹, Valentina Lara-Erazo, MD², Sabry Ayad, MD¹, Arthur Calimaran, MD³, Monica Cheriyan, MD, MPH³.
 Department of Anesthesiology¹, Department of Outcomes Research², Department of Obstetric Anesthesiology³
 Cleveland Clinic Foundation, Cleveland, OH

Introduction

- ❖ Pregnancy and obesity are high risk factors of severe COVID 19
- ❖ Long term outcomes include residual pulmonary fibrosis
- ❖ Risk of placental and fetal hypoxia

Case Description

- ❖ 22-year-old 37w4d primigravida
- ❖ Cesarean section indication: morbid obesity and pulmonary fibrosis.
- ❖ Pathologies:
 - ❖ Pulmonary fibrosis: Severe COVID 19 infection one year before: 40 days mechanical ventilation and tracheostomy (Fig A-D)
 - ❖ Tachycardia
 - ❖ Morbid Obesity (BMI 45.7)
- ❖ Pharmacological therapy:
 - ❖ Rate control medication
 - ❖ Home oxygen therapy through nasal cannula at 3L/minutes to maintain a SpO2 of 95%
- ❖ Prenatal care: Dyspnea and palpitations with increased activities. No history of sleep apnea, but endorsed using a couple of pillows at night.
 - ❖ Transthoracic echocardiogram within normal ranges




Fig A. Intubation 10/14/2021




Fig B. 11/19/2021




Fig C. 11/24/2021 tracheostomy





Fig D. 12/21/2021 Tracheostomy out



CESAREAN SECTION WITH EPIDURAL ANESTHESIA IN A MORBIDLY OBESE PARTURIENT WITH POST-COVID PULMONARY FIBROSIS

Yuta Kashiwagi, MD¹, Valentina Lara-Erazo, MD², Sabry Ayad, MD¹, Arthur Calimaran, MD³, Monica Cheriyan, MD, MPH³.
Department of Anesthesiology¹, Department of Outcomes Research², Department of Obstetric Anesthesiology³
Cleveland Clinic Foundation, Cleveland, OH

Case Description (continued)

Physical Exams:

- ❖ Height 180.3cm, weight 148.6kg, BMI 45.7
- ❖ Blood pressure 110/74mmHg, Pulse 118 bpm, sinus rhythm
- ❖ SpO2 95% with oxygen through nasal cannula at 3L/minutes
- ❖ Head and neck: thick and short neck, anterior cervical scar of pervious tracheostomy, no difficulty swallowing, no voice changes, no obstructive sensation while breathing.

Anesthesia management

- ❖ Epidural anesthesia
 - ❖ T12/L1 level with a needle depth of 7cm
 - ❖ Catheter left 13 cm from the skin
 - ❖ Four doses of 100mg 2% lidocaine with 25mcg epinephrine
 - ❖ T5 level bilateral block
 - ❖ Blood pressure was maintained with continuous phenylephrine infusion with intermittent boluses.
- ❖ Discharged without complications two days later.

Discussion

- ❖ Cesarean delivery early in the third trimester to avoid a further decrease in lung capacity at the moment of labor
- ❖ Anesthesia care
 - ❖ High block levels could deteriorate respiratory function and be harmful for the fetus.
 - ❖ Avoiding Radical sympathectomy and leading tachycardia
 - ❖ Challenging for general anesthesia conversion
 - ❖ Respiratory insufficiency
 - ❖ Unfavorable airway
- ❖ Alternative: Combined Spinal Epidural (CSE) could have been another option.

Conclusion

- ❖ Cesarean section in a morbidly obese parturient with post-COVID pulmonary fibrosis was successfully managed with epidural anesthesia.