

Unexpected prolonged neuromuscular blockade in an eclamptic patient requiring emergent caesarean delivery

Kathryn Clark MB BCh BAO, Giselle Villar MD FRCPC, Anthony Chau MD FRCPC



Case presentation:

- 31 y.o. (72.3 kg) G1P0 at 38+4 GA with severe hypertension, headache and blurry vision
- Plan: Magnesium, nifedipine + urgent caesarean delivery
- Prior to OR → seizure + prolonged fetal bradycardia
- RSI (propofol + succinylcholine 160 mg) → emergent delivery of fetus
 - Maintenance: propofol, remifentanyl, sevoflurane
- Persistent motor blockade after cessation of anesthesia x 20 minutes

Two hours after initial succinylcholine administration

- Evidence of muscle blockade with peripheral nerve stimulation
 - Neuroimaging with CT head

Three hours after initial succinylcholine administration


- Extubation criteria met and endotracheal tube removed

Table 1. *Laboratory results*

Platelets	75 x 10 ⁹ /L
Urine P:Cr	2859 g/mol
Bilirubin	55 umol/L (<2)
AST	446 U/L (<40)
ALT	342 U/L (<55)
LDH	1424 U/L (<250)

Unexpected prolonged neuromuscular blockade in an eclamptic patient requiring emergent caesarean delivery

Kathryn Clark MB BCh BAO, Giselle Villar MD FRCPC, Anthony Chau MD FRCPC



Differential Diagnosis:

- Pseudocholinesterase Deficiency
 - PChE activity measured (collected 1d PP)
 - Cholinesterase, total 192 U/L (620-1370 U/L)
 - Cholinesterase phenotype: homozygous for typical enzyme
 - ? reduced activity due to pregnancy and liver dysfunction
- Magnesium + Nifedipine
 - Magnesium level 1.91 mmol/L 5 hours following delivery
 - ? Potentiation of NMB from magnesium
- Intracranial Process
 - CT and MRI revealed changes consistent with posterior reversible encephalopathy syndrome (PRES)

Phenotype	Dibucaine #
Homozygous typical	70-80
Heterozygous atypical	50-60
Homozygous atypical	20-30

Unexpected prolonged neuromuscular blockade in an eclamptic patient requiring emergent caesarean delivery

Kathryn Clark MB BCh BAO, Giselle Villar MD FRCPC, Anthony Chau MD FRCPC



Discussion:

- Multiple differential diagnoses in a parturient with prolonged neuromuscular blockade
 - PChE activity can be evaluated by studies which investigate activity level and gene phenotype

- Repeat testing can help reveal the extent to which pregnancy and liver disease can affect cholinesterase activity
 - This should be completed >2 months post-delivery after activity levels are expected to normalize

- Controversy whether routine peripheral nerve monitoring of neuromuscular blockade should be mandatory following succinylcholine
 - Residual blockade confirmed with peripheral nerve stimulation + allowed for continued monitoring of resolution of block