


Anesthetic management of obstetric patient with large parieto-occipital meningioma: a case report

Rebecca Thompson, MD; Kristin Falce, MD; Barbara Orlando, MD, Ana Lisa Ramirez-Chapman, MD

- Incidence of brain tumors in pregnancy is similar to nonpregnant women of the same age
- Previously asymptomatic neoplasms may enlarge during pregnancy and become symptomatic with multiple potential implications for the anesthesiologist


Case: 40-year-old parturient presented in her third trimester with 10-day history of worsening headache, visual changes, memory dysfunction, gait disturbance



MRI brain, sagittal and axial images of 5cm intracranial mass

Anesthetic management of obstetric patient with large parieto-occipital meningioma: a case report

- Cesarean section under general anesthesia
 - 2/2 elevated ICP
 - Pre-induction radial arterial line
 - Titrated phenylephrine and nicardipine PRN for MAP goals > 70
 - 2 large bore PIVs
 - Induction meds: propofol 200mg, rocuronium 100mg, remifentanyl gtt 0.2 mcg/kg/min
 - Post-op bilateral quadratus lumborum blocks w/ 0.25% bupivacaine prior to extubation
- Delivery of healthy baby w/o complications: APGARs 1, 6, 8
- 24 hours post cesarean → Craniotomy for mass resection
- Mom and baby doing well to date; mom 90% back to baseline



Anesthetic management of obstetric patient with large parieto-occipital meningioma: a case report

CONCLUSION

- Space-occupying intracranial lesions in pregnant women; implications for delivery:
 - Higher likelihood of cesarean delivery with general anesthesia
 - Decrease risk of increased ICP during labor
 - Risk of herniation from neuraxial anesthesia
- These cases require more communication, consideration, and planning on the part of the obstetric anesthesiologist
- Imperative to seek advice from neurologists and neurosurgeons in each case, as there is insufficient data to make general recommendations regarding delivery and anesthesia plans

