

Management of an Urgent Cesarean Section in a Patient with Unknown Severe Pulmonary Hypertension, Obesity, Heart Failure with Reduced Ejection Fraction and Prior Classical Cesarean Section

Tyler Guidugli, DO; Tayler Hines, DO; Jon Holzberger, MD; Johannes Steyn, MD; Regina Fragneto, MD; James Damron, MD
University of Kentucky Department of Anesthesiology, Lexington, KY



Hospital Presentation

- 33 year old G8P1 at 31 weeks transfer for vaginal bleeding and contractions. In the days leading up to her presentation she experienced new onset angina, headache, and vision changes.
- Past medical history: class 3 obesity (BMI 82), HFrEF, COPD on 2 liters home oxygen, HTN, prior classical cesarean delivery, multiple prior MI and PE requiring therapeutic anticoagulation, and limited prenatal care.
- On arrival, she was requiring 5 liters nasal cannula and started on tocolytics while a cardiopulmonary work up was performed. Prior transthoracic ECHO in 2017 revealed LVEF 55%.



Anesthetic Course

Pre-op

- General anesthesia chosen due to the patient being on therapeutic enoxaparin
- Transthoracic ECHO revealed LVEF 30-35% and RVSP 50-70mmHg
- Chest x-ray revealed enlarged cardiac silhouette
- Pre-induction arterial line and post induction central line was placed as well as large bore peripheral access

Intra-op

- TEE placed revealing severe biventricular dysfunction, severe pulmonary hypertension and severe TR
- Patient started on epinephrine, norepinephrine, vasopressin and milrinone due to TEE findings
- She stabilized after inotropic support and baby was delivered uneventfully
- She was then transferred to the MICU intubated

Post-op

- She became increasingly unstable with worsening oxygenation ultimately requiring peripheral VA ECMO
- POD 1 her arterial cannula became dislodged resulting in a ACLS
- After LV recovery an attempt was made to transition to an RVAD, unfortunately she was unable to wean from RVAD support and was transitioned to comfort care



Figure 1: Intra-op
Mid-esophageal 4
chamber TEE




Figure 2: Dilated PA

Management of an Urgent Cesarean Section in a Patient with Unknown Severe Pulmonary Hypertension, Obesity, Heart Failure with Reduced Ejection Fraction and Prior Classical Cesarean Section

Tyler Guidugli, DO; Tayler Hines, DO; Jon Holzberger, MD; Johannes Steyn, MD; Regina Fragneto, MD; James Damron, MD
University of Kentucky Department of Anesthesiology, Lexington, KY



Discussion

- Cardiovascular disease is now the leading cause of maternal mortality in developed nations
- Pulmonary hypertension in concurrence with HFrEF has a significantly higher morbidity and mortality in pregnancy than pulmonary hypertension alone
- Due to the urgency of the case, her late presentation and limited prenatal care, we had no opportunity for optimization prior to delivery
- The use of transesophageal ECHO greatly impacted our management by earlier identification of the severity of right heart failure and implementation of inotropic agents
- Epidural is the preferred modality for cesarean delivery in patients with significant cardiovascular conditions

References

1. Pieper PG, Hoendermis ES. Pregnancy in women with pulmonary hypertension. *Neth Heart J*. 2011 Dec;19(12):504-8. doi: 10.1007/s12471-011-0219-9. PMID: 22068738; PMCID: PMC3221745.
2. Thomas E, Yang J, Xu J, Lima FV, Stergiopoulos K. Pulmonary Hypertension and Pregnancy Outcomes: Insights From the National Inpatient Sample. *J Am Heart Assoc*. 2017 Oct 24;6(10):e006144. doi: 10.1161/JAHA.117.006144. PMID: 29066439; PMCID: PMC5721838.

