



Ischemic Cardiomyopathy Discovered During Pregnancy

Jessie Merrill, MD & Kasey Fiorini, MD



THE OHIO STATE UNIVERSITY

Case Presentation

24w6d

Heparin gtt for anticoagulation. Labetalol/hydralzine for BP/HFrEF. Insulin gtt for hyperglycemia (BG 200-300s). Suboxone for opioid withdrawal. IV furosemide for diuresis.

26w0d-26w3d

Persistent **delirium with c/f cyanide toxicity**. Persistent cardiogenic shock. Not candidate for advanced mechanical therapies given substance abuse.

27w0d

Repeat LTCS with BTL under GETA with intraop TEE, PAC and arterial line. Remained intubated postop.

Discharge & Beyond

Left AMA POD 12. Minimal follow-up. Continues to be on GDMT with life vest.

24w5d

43 year old G3P1 with PSA (meth, opioid) presented with CP, SOB, severe HTN and **left CVA** symptoms. TTE with **EF 20-25%**, global hypokinesis and **LV thrombus**. EKG with minimal ST changes V2/V3. Normal troponin. CT head with left parietal, frontal and occipital lobes. **Gross volume overload** on exam.

25w3d-26w0


Polymorphic NSVT. Started on lidocaine gtt. After multidisciplinary conversation, LHC/RHC showed **severe CAD** (mLAD 100%, RCA 100%, D1 90%) and **cardiogenic shock** (elevated filling pressure, low cardiac index). Nitroprusside gtt for afterload reduction. Furosemide gtt started.

26w5d

Repeat RHC with PAC left in situ. Persistent volume overload and cardiogenic shock. **Milrinone gtt** for inotropic support (previous held due to risk of ventricular ectopy). Able to wean nitroprusside with improvement in delirium.

POD 1-3

Extubated POD1. Started on GDMT. POD3 all vasoactive meds weaned. No longer in cardiogenic shock. **No further decompensation postpartum**. MRSA bacteremia.



Discussion Points

For patients with undifferentiated peripartum heart failure, one must consider ischemic cardiomyopathy as possible etiology, especially in patients with multiple risk factors.

- **RF:** tobacco use, DM, HTN, meth/cocaine use, obesity, AMA, hypertriglyceridemia, family history of MI <60 years old
- **Labs/Studies:** q waves/ST changes on EKG, regional wall motion changes, elevated troponin, polymorphic VT
- **Prevalence:** 1:10,000 (ischemic heart disease) 7.5:100,000 (risk of MI in pregnancy)¹

Physiologic changes of pregnancy can exacerbate underlying cardiac dysfunction.

- **Increased myocardial O₂ demand:** ↑ cardiac output ↑ circulating blood volume ↑ oxygen consumption ↑ HR
- maximally increased ~ 34 weeks

Pregnant patients in cardiogenic shock require careful multidisciplinary management.

- **Delivery planning:**
 - **Key Players:** cardiology (EP, heart failure, cardio-obstetrics), CT surgery, MFM, obstetric & cardiac anesthesia, NICU, ICU & L&D nursing
 - **Key Decisions:** mode & timing of delivery, location of delivery and postop recovery, hemodynamic monitoring (TEE, PAC, arterial line, cardiac output)
- **Early discussion of candidacy for advanced therapies:** ECMO, LVAD, heart transplant

For patients with ischemic cardiomyopathy, the postpartum period deserves equal consideration.

- **Highest risk of cardiac decompensation:** ↑ cardiac output ↑ SVR
- **Revascularization:** PCI considered safe during pregnancy, CABG requiring CPB usually deferred until post partum given high risk of fetal demise (20-40%)²
- **Breast-feeding:** ARNi (*Entresto*: sacubitril/valsartan) contraindicated, most GDMT safe for lactation but may require special monitoring of infant
- **Family planning**

¹Davies GAL, Herbert WNP. Ischemic heart disease and cardiomyopathy in pregnancy. J Obstet Gynaecol Can. 2007 Jul;29(7):575-579

²van Steenberg GJ, Tsang QHY, van der Heijden OWH, Vart P, Rodwell L, Roos-Hesselink JW, van Kimmenade RRJ, Li WWL, Verhagen AFTM. Timing of cardiac surgery during pregnancy: a patient-level meta-analysis. Eur Heart J. 2022 Aug 1;43(29):2801-2811

