


Introduction

- Peripartum Cardiomyopathy (PPCM) is a type of non-ischemic cardiomyopathy with reduced ejection fraction affecting a parturient in late pregnancy or in the first five months postpartum
 - Risk factors include hypertension, preeclampsia, age >30 years, African American race, multigestational pregnancy
- Diagnosis is often difficult and may be delayed, with many symptoms similar to those seen in normal pregnancy
 - Symptoms: dyspnea, orthopnea, chest tightness
 - Physical exam: tachycardia, tachypnea, edema, elevated jugular venous pressure
 - Echocardiography: LVEF <45%, possible LA or biatrial enlargement
 - Laboratory findings: elevated BNP
 - Chest X-ray: pulmonary venous congestion
- Management of PPCM is comparable to that of other forms of heart failure, with severe cases sometimes requiring advanced therapies
 - Mechanical circulatory support
 - Heart transplantation



Getting to the Heart of the Matter: A Case of Peripartum Cardiomyopathy Leading to Heart Transplantation

Domenic Pedulla MD¹, Kristine Villegas MD¹, Christina Zecca, MD¹, Danielle McCullough MD¹, Sharon Abramovitz MD¹

¹Department of Anesthesiology, Weill Cornell Medical College, New York, NY

Case	<ul style="list-style-type: none">34 year old female, G1P0, with PMSH anemia, leiomyoma, prior cholecystectomyPresented at 37w5d with chest pain, dyspnea, non-productive cough
Initial Work up	<ul style="list-style-type: none">Elevated BNP, Negative troponin, Negative CT PEBedside TTE with severely reduced LVEF (17%) and moderately reduced RV functionMultidisciplinary team including Obstetric Anesthesia, Cardiothoracic Surgery, Heart Failure Cardiology, OB/GYN assembled
Preoperative Optimization	<ul style="list-style-type: none">Arterial line placed, inotropic support with dobutamine initiated pre-operatively
Emergent Cesarean Delivery	<ul style="list-style-type: none">General anesthesiaCentral access obtained, Swan-Ganz catheter placedEpinephrine infusion initiatedIntra-aortic balloon pump placedCCU transfer
Postoperative Course	<ul style="list-style-type: none">PPD1- Worsening biventricular functionPPD2- VA ECMO initiated with peripheral ImpellaPPD8- Transferred to an affiliated hospital for cardiac transplant evaluationPPD9- Developed paraplegia and saddle anesthesia, diagnosed with spinal cord infarction, failed trial of decannulationPPD 11- Urgent chest exploration with concern for tamponade
UNOS Status 1 for Transplant on PPD13	<ul style="list-style-type: none">PPD17- Successfully underwent orthotopic heart transplantDischarged to spinal cord rehabilitation center
Long Term Follow Up	<ul style="list-style-type: none">Follow up TTE 2 months postoperatively demonstrated LVEF 65% with mildly reduced RV function




Figure 1. Peripartum Vital Signs and Medication Timeline

Getting to the Heart of the Matter: A Case of Peripartum Cardiomyopathy Leading to Heart Transplantation

Domenic Pedulla MD¹, Kristine Villegas MD¹, Christina Zecca, MD¹, Danielle McCullough MD¹, Sharon Abramovitz MD¹

¹Department of Anesthesiology, Weill Cornell Medical College, New York, NY



Discussion

- Vaginal delivery is preferred in cases of PPCM, but both neuraxial and general anesthesia can be safely performed when cesarean delivery is required
 - Focus on strict hemodynamic control with careful cardiac monitoring
- Period immediately following delivery can be particularly tenuous with acute increases in venous return increasing risk of fluid overload
- Acutely decompensated patients should be managed by specialized multidisciplinary teams
 - Treatment may include vasodilators, diuretics, and inotropic support
 - Mechanical circulatory support with an intra-aortic balloon pump, VA ECMO, or ventricular assist devices should be considered early in refractory cases
- Approximately 10% of PPCM patients require heart transplantation
 - Data suggest higher rates of graft failure and lower age-adjusted survival compared to non-PPCM patients
- Long-term follow up is required, though questions remain about optimal duration of medication treatment after recovery

References:

- 1- Davis M.B., et al. Peripartum Cardiomyopathy: JACC State-of-the-Art Review. J Am Coll Cardiol. 2020 Jan 21;75(2):207-221.
- 2- Kwon, J., et al. Heart Transplantation for Peripartum Cardiomyopathy: Outcomes Over 3 Decades. Ann Thorac Surg. 2022 Sept; 114(3):650-658.
- 3- Ramachandran R, et al. Anesthetic management of patients with peripartum cardiomyopathy. J Obstet Anaesth Crit Care. 2011; 1:5-12.

Questions/Comments: Please contact Domenic Pedulla MD: djp9016@nyp.org