




Point of care strain analysis during vaginal delivery and third stage of labor: A case series

M. Smith, M. Montes, B. Scavone, D. Tobes, N. Cole, S. Shahul, C. Thomas


- Emerging evidence that parturients with hypertensive diseases of pregnancy exhibit myocardial strain during early labor¹
- Signs of potential myocardial injury have been identified in at delivery in otherwise healthy parturients²




Pos 3: Parasternal LV short axis



Pos 2: Apical 4-chamber



Pos 3: Parasternal long axis



Pos 2: Apical 5 - Chamber

Early 2nd
stage

Maternal
Pushing

Post-
Infant
delivery

Post-
Placental
Delivery



AT THE FOREFRONT
UChicago Medicine

SOAP
Society for Obstetric
Anesthesia and Perinatology


Case Series



1

37yo G1P0 41w
w/ neuraxial

- ✓ **Analyzable images at all time-points**
 - Segmental strain analysis **abnormal** in 3 segments post-fetal delivery
 - Long second stage, 2 hours of maternal pushing + forceps delivery



2

30yo G2P1 39w
w/ neuraxial

- ✓ Analyzable images **for speckle tracking only; poor quality A4+5C**
 - Imaging during maternal pushing limited: **delivered on hands/knees**
 - No abnormal strain



3

30yo G8P6016 36w
w/ neuraxial

- ✓ **Analyzable images at two time-points**
 - Segmental strain analysis and diastology was **abnormal** in one segment during delivery
 - Precipitous delivery: only 3 full sets of images obtained



AT THE FOREFRONT
UChicago Medicine

SOAP
Society for Obstetric
Anesthesia and Perinatology

Discussion

- **Novel use** of serial TTE exams during parturition
- Strain analysis during vaginal delivery **is feasible**
- Limited by positioning and rapid progression of labor
- Future directions:
 - Characterize strain patterns in women with hypertensive diseases of pregnancy
 - Compare strain patterns **within subjects** at different time points
 - Determine normative strain values

