Efficacy and Safety Is Maintained in Adult Patients With Paroxysmal Nocturnal Hemoglobinuria Receiving Pegcetacoplan for Up to 3 Years

MDS-397

Carlos de Castro[1]; Brian Mulherin[2,3]; Christopher J. Patriquin[4]; Veena Selvaratnam[5]; Raymond Siu Ming Wong[6]; Richard J. Kelly[7]; Lisa Tan[8,9]; Carl Friddle[10]; Peter Hillmen[10]; Regis Peffault de Latour[11,12]

[1]Duke University, Durham, NC, USA; [2]Hematology Oncology of Indiana, Indianapolis, IN, USA; [3]Ascension St. Vincent Carmel, Carmel, IN, USA; [4]University Health Network, Toronto, ON, Canada; [5]Ampang Hospital, Ampang, Malaysia; [6]The Chinese University of Hong Kong, Prince of Wales Hospital, Hong Kong, Hong Kong; [7]Department of Haematology, St. James's University Hospital, Leeds, UK; [8]Swedish Orphan Biovitrum AB, Stockholm, Sweden; [9]Lisa Tan Pharma Consulting Ltd., Cambridge, UK; [10]Apellis Pharmaceuticals, Inc., Waltham, MA, USA; [11]French Reference Center for Aplastic Anemia and Paroxysmal Nocturnal Hemoglobinuria, Paris, France; [12]Université Paris Cité, Paris, France

CONCLUSIONS

- ✓Pegcetacoplan is the first proximal inhibitor demonstrating long-term efficacy and safety for a broad PNH population
- ✓Hematologic improvements were maintained through 3 years in C5i-experienced patients (PEGASUS) and through 2.5 years in C5i-naive patients (PRINCE)
- ✓Transfusion burden was markedly reduced after pegcetacoplan initiation, with 52% of patients avoiding transfusion through 3 years in PEGASUS and 67% through 2.5 years in PRINCE
- ✓No new safety findings were identified

INTRODUCTION

- Pegcetacoplan increased Hb concentrations in C5i-experienced and -naive adults with PNH in 2 phase 3 clinical trials (PEGASUS [NCT03500549] and PRINCE [NCT04085601], respectively)[1–3]
- Long-term safety and efficacy of pegcetacoplan are being assessed in the ongoing 307 OLE (NCT03531255)

OBJECTIVE

To report the longest continuous clinical trial data set for pegcetacoplan by conducting an integrated analysis of 2 phase 3 trials and the subsequent OLE

METHODS


- Patients in the 307 OLE initially received pegcetacoplan 1080 mg subcutaneously twice weekly (dose escalations allowed)
- Efficacy was assessed from pegcetacoplan initiation up to 3 years (PEGASUS) and 2.5 years (PRINCE) by Hb, LDH, ARC, and indirect bilirubin values; FACIT-Fatigue scores; and transfusion avoidance rates
  - The Hb analysis accounted for transfusions; measures within 28 days before pegcetacoplan initiation and within 60 days of transfusion during pegcetacoplan treatment were excluded
- Safety during pegcetacoplan monotherapy was assessed

RESULTS

- Of 133 patients in the phase 3 trials, 114 enrolled in the 307 OLE (PEGASUS, 64; PRINCE, 50)
- In the year before initial enrollment, ≥75% received a transfusion
- In PEGASUS and PRINCE, prior to pegcetacoplan initiation, mean (SD) Hb concentrations were 9.0 (1.1) g/dL and 9.3 (1.4) g/dL, respectively; median (IQR) LDH concentrations were 217 (185, 277) U/L and 1964 (1409, 2503) U/L; mean (SD) FACIT-Fatigue scores were 31.6 (11.7) and 36.6 (10.0); mean (SD) ARCs were 214 (79) ×10^9 cells/L and 181 (69) ×10^9 cells/L; mean (SD) indirect bilirubin concentrations were 33.2 (24.9) µmol/L and 32.3 (16.7) µmol/L


RESULTS (*continued*)

Figure 1. Hb concentrations through up to 3 years of pegcetacoplan treatment[a]




[a]Patients who had a transfusion were not included in the Hb analysis for 60 days after the transfusion. [b]Normal ranges are per central laboratory; local laboratory values were normalized using central laboratory normal ranges.
BL, baseline (at pegcetacoplan initiation); Hb, hemoglobin; LLN, lower limit of normal (12 g/dL for female patients and 13.6 g/dL for male patients); SE, standard error.

Figure 2. LDH concentrations through up to 3 years of pegcetacoplan treatment



[a]Normal ranges are per central laboratory; local laboratory values were normalized using central laboratory normal ranges.
BL, baseline (at pegcetacoplan initiation); IQR, interquartile range; LDH, lactate dehydrogenase; ULN, upper limit of normal (226 U/L).


Figure 3. FACIT-Fatigue scores through up to 3 years of pegcetacoplan treatment[a]



[a]Greater FACIT-Fatigue scores indicate less fatigue. [b]Reported by Cella and colleagues.[4]
BL, baseline (at pegcetacoplan initiation); FACIT-Fatigue, Functional Assessment of Chronic Illness Therapy-Fatigue; SE, standard error.


- With pegcetacoplan, Hb (**Figure 1**) and LDH (**Figure 2**) concentrations markedly improved and remained stable through up to 3 years
- FACIT-Fatigue scores rapidly increased (indicating less fatigue) and were maintained near the general population norm of 43.6[4] (**Figure 3**)

Figure 4. ARCs through up to 3 years of pegcetacoplan treatment




[a]Normal ranges are per central laboratory; local laboratory values were normalized using central laboratory normal ranges.
ARC, absolute reticulocyte count; BL, baseline (at pegcetacoplan initiation); SE, standard error; ULN, upper limit of normal (120x10^9 cells/L).

Figure 5. Indirect bilirubin levels through up to 3 years of pegcetacoplan treatment



[a]Normal ranges are per central laboratory; local laboratory values were normalized using central laboratory normal ranges.
BL, baseline (at pegcetacoplan initiation); SE, standard error; ULN, upper limit of normal (15.4 µmol/L).

Figure 6. Transfusion avoidance through up to 3 years of pegcetacoplan treatment



- ARC improvements were maintained below the ULN (**Figure 4**), as were indirect bilirubin concentrations (**Figure 5**)
- Annual transfusion avoidance ranged from 71.2%–79.2% (PEGASUS) to 79.5%–86.4% (PRINCE) (**Figure 6**)
- In the PEGASUS population, 51.6% avoided transfusion for up to 3 years; in the PRINCE population, 66.5% avoided transfusion for up to 2.5 years (**Figure 6**)

Table. TEAEs during pegcetacoplan monotherapy for up to 3 years

Patients with events, n (%)	PEGASUS N=80	PRINCE N=52	Total N=132
TEAEs	79 (98.8)	50 (96.2)	129 (97.7)
Considered related to pegcetacoplan	43 (53.8)	17 (32.7)	60 (45.5)
Serious TEAEs	43 (53.8)	23 (44.2)	66 (50.0)
Considered related to pegcetacoplan	5 (6.3)	1 (1.9)	6 (4.5)
TEAEs leading to pegcetacoplan discontinuation	16 (20.0)	1 (1.9)	17 (12.9)
TEAEs leading to death	1 (1.3)	3 (5.8)	4 (3.0)
Considered related to pegcetacoplan	0	0	0
TEAEs of special interest	70 (87.5)	46 (88.5)	116 (87.9)
Hypersensitivity	27 (33.8)	16 (30.8)	43 (32.6)
Infections	63 (78.8)	35 (67.3)	98 (74.2)
Serious infections	17 (21.3)	10 (19.2)	27 (20.5)
Sepsis	4 (5.0)	2 (3.8)	6 (4.5)
Meningitis	0	0	0
Thrombosis	3 (3.8)	0	3 (2.3)
Breakthrough hemolysis[a]	23 (28.8)	16 (30.8)	39 (29.5)
Patients with 1/2 events	14 (17.5) / 4 (5.0)	10 (19.2) / 3 (5.8)	24 (18.2) / 7 (5.3)

[a]Clinically significant (as reported by investigators) and laboratory-confirmed events. AE, adverse event; TEAE, treatment-emergent adverse event (AEs with a start date on or after the first dose of pegcetacoplan monotherapy or with a start date before the date of the first dose of pegcetacoplan but increased in severity on or after the date of the first dose of drug. Did not include AEs occurring more than 8 weeks after the last dose of pegcetacoplan). Patients with multiple occurrences of an AE were counted only once.

- Most patients had a TEAE during up to 3 years of pegcetacoplan monotherapy (**Table**)
- Serious TEAEs occurred in 50.0% of patients overall, and 4.5% of all patients had a serious TEAE deemed pegcetacoplan related
- Overall, 17 patients discontinued pegcetacoplan due to a TEAE; of those, 9 discontinued due to a hemolytic disorder, with 7 of these discontinuing within 1 year of starting pegcetacoplan
- In all, 4 (3.0%) TEAE-related deaths occurred (none deemed pegcetacoplan related) and 3 (2.3%) patients had a thrombotic event
- Breakthrough hemolysis occurred in 28.8% (23 of 80) of patients in PEGASUS and 30.8% (16 of 52) of patients in PRINCE
- No meningitis cases were reported

REFERENCES

1. Hillmen P, et al. *N Engl J Med*. 2021;384(11):1028–37.
2. de Latour RP, et al. *Lancet Haematol*. 2022;9(9):e648–59.
3. Wong RSM, et al. *Blood Adv*. 2023;7(11):2468–78.
4. Cella D, et al. *Cancer*. 2002;94(2):528–38.

Abbreviations:
ARC, absolute reticulocyte count; BL, baseline; C5i, complement component 5 inhibitor; FACIT-Fatigue, Functional Assessment of Chronic Illness Therapy-Fatigue; Hb, hemoglobin; IQR, interquartile range; LDH, lactate dehydrogenase; LLN, lower limit of normal; OLE, open-label extension; PNH, paroxysmal nocturnal hemoglobinuria; SD, standard deviation; SE, standard error; TEAE, treatment-emergent adverse event; ULN, upper limit of normal.

12th Annual Meeting of the Society of Hematologic Oncology (SOHO) | September 4–7, 2024 | Houston, TX
Prior presentation: 65th Annual Meeting and Exposition of the American Society of Hematology (ASH); December 9–12, 2023; San Diego, CA

Disclosures: **CdC** reports consultancy at Apellis; data safety monitoring board at Regeneron; advisory board at Omeros; honoraria from Novartis, Alexion, BioCryst, and Apellis; and research funding from Alexion and Apellis. **BM** has nothing to disclose. **CJP** has received consultant fees from Alexion, Apellis, Regeneron, Takeda, BioCryst, Sobi, Amgen, and Novartis. **VS** has received consultant fees from Apellis. **RSMW** has received consulting fees, honoraria, research funding, and speaker's bureau fees from Alexion and F. Hoffmann-La Roche Ltd.; and research funding and speaker's bureau fees from Apellis. **RJK** has received research funding from Novartis and consulting fees, honoraria, and speaker's bureau fees from Novartis, Roche, Alexion, Sobi, Jazz, Pfizer, Astellas, Amgen, AbbVie, and Biologix. **LT** is a consultant for SOBI. **PH**, and **CF** are employees of Apellis and hold stock options. **RPdL** has nothing to disclose.
The authors also thank Dale Zhang and Jessica Savage for their input on early versions of this presentation.

Acknowledgments: Swedish Orphan Biovitrum AB and Apellis Pharmaceuticals, Inc., provided funding toward this study. Medical writing support was provided by Catherine Champagne, PhD, and Kathryn Fogarty, PhD, both of Kay Square Scientific, funded by Apellis Pharmaceuticals, Inc., and Swedish Orphan Biovitrum AB.