# Implementing Prefilled Sterile Syringes in the Pediatric Emergency Department of a Tertiary Care Hospital, Pakistan: A Novel Approach to Enhance Medication Safety

Authors: Parveen, Fezzia J. | Senior Pharmacist, Saba, Javed J | Pharmacist
Aga Khan University Hospital, Karachi, Pakistan

## Introduction

Pediatrics are a vulnerable group for dosing and medication errors. Most of the drugs used are available in dosages and units ready for dispensing in adults. This necessitates a lot of calculations and has the potential for errors.

## Objectives

The aim of this study was to evaluate the efficacy of Implementing Prefilled Sterile Syringes in the Pediatric (PPSs) in reducing time of medication administration to critically ill pediatric patients in the emergency department (ED) of a tertiary care hospital in Karachi, Pakistan.

## Methods

This was a quasi-experimental study done on the quality project from January 2018 to March 2018 in the ED of a tertiary care hospital in Karachi, Pakistan. This project was implemented by adopting Plan-Do-Study-Act (PDSA) methodology. In this study, mean reduction in intravenous medication administration time was measured. The most frequently used pediatric emergency medications were selected in particular strengths for the pilot as ready-to-administer PPSs: magnesium sulfate 5 meq /2 mL, 3% hypertonic saline 150 mL piggyback , and paracetamol 200 mg/20 mL. Pre- and postintervention data were collected and analyzed inferentially in SPSS.

**Figure 2 Prefilled syringes implementing the Pediatric Emergency Department**




Figure 2: Prefilled syringes implementing the Pediatric Emergency Department

## Results

We observed a mean reduction in dose administration time from 156 to 94.5 seconds (95% CI, 111.9–129.4) for magnesium sulfate 5 meq /2 mL and from 304 to 41 seconds (95% CI, 251.1–356.9) for hypertonic saline 150 mL. Inferentially, the mean reduction in time was significant (P<0.01) after implementing both PPSs.




## Summary

Implementing ready-to-administer PPSs in pediatric emergency patients not only significantly reduces the medication administration time but is also a good strategy to enhance patient safety in pediatric population. It has the potential to improve throughput in the ED.

## Discussion

Our study highlighted that needs to administer prefilled sterile syringes has significantly reduce the time of medication administration in pediatric patients. Reduction in medication administration time has the potential for improving process flow in busy emergency department. The PPSs also have the potential to improve patient safety by reducing the number of steps and reliance than preparation. Additionally, it also reduces unnecessary calls for clarifications and queries about medication orders.[1]

The dose of magnesium sulfate was 50 mg/kg, and 3% hypertonic saline was 5 ml/kg as calculated in this project, as the dose was 100 mg/kg reduced from 500 mL to 150 mL. This adjustment was made primarily for safety reasons and to avoid fluid overload, which can lead to serious adverse events, associated with the high concentration electrolytes.

## Key words

Prefilled sterile syringes, Pediatric patients, Emergency Department, Medication Error, Patient Safety

## References

1. Israel T, Waddick A, Hollenbeck J, Avery L, Pichel J, Gregorio C, et al. Using Pre-Filled Medication Safety Self-Assessment™ to improve medication use processes. The Joint Commission Journal on Quality and Safety. 1998;29(6):311–26.



SOHO2025
Poster Number: CT-192