

Correlation Between IL-8 and TNF-Alpha and Overall Survival in Imetelstat-Treated Patients With Myelofibrosis Relapsed or Refractory to a Janus-Associated Kinase Inhibitor in the IMbark Trial

John O. Mascarenhas, MD,¹ Claire N. Harrison, MD,² Alessandro Lucchesi, MD, PhD,³ Angela Fleischman, MD, PhD,⁴ Rami S. Komrokji, MD,⁵ Alessandro M. Vannucchi, MD,⁶ Jean-Jacques Kiladjian, MD, PhD,⁷ Yiqun Yang, PhD,⁸ Libo Sun, PhD,⁸ Michelle Mudge-Riley, DO, MHA,⁸ Fei Huang, PhD,⁸ Prithviraj Bose, MD⁹

¹Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA; ²Guy's and St Thomas' National Health Service Foundation Trust, London, UK; ³Hematology Unit, IRCCS Istituto Romagnolo per lo Studio dei Tumori (IRST) "Dino Amadori", Meldola, Italy; ⁴University of California, Irvine, Irvine, CA, USA; ⁵Moffitt Cancer Center, Tampa, FL, USA;

⁶University of Florence, Florence, Italy; ⁷Hôpital Saint-Louis, Assistance Publique-Hôpitaux de Paris, Université Paris Cité, Paris, France; ⁸Geron Corporation, Foster City, CA, USA; ⁹The University of Texas MD Anderson Cancer Center, Houston, TX, USA

MPN - 298



Copies of this poster obtained through Quick Response (QR) Code are for personal use only and may not be reproduced without permission

Background


- The dysregulation of inflammatory cytokines plays a key role in the pathogenesis, progression, and clinical features of myelofibrosis (MF)^{1,2}
- In particular, increased interleukin (IL)-8 and tumor necrosis factor alpha (TNF-α) have been associated with poor prognosis in patients with MF^{1,3}
- In preclinical evaluations, treatment with the telomerase inhibitor, imetelstat, led to decreased expression of IL-8 in various tumor cell lines⁴
- In the Phase 2 IMbark trial (NCT02426086) of 2 dosing regimens, treatment with imetelstat showed a dose-related trend for clinical benefit, with improved overall survival (OS) in the higher dose arm compared with the lower dose arm, an acceptable safety profile, and potential disease-modifying activity in patients with intermediate-2-risk or high-risk MF relapsed or refractory (R/R) to Janus kinase inhibitor (JAKi) treatment^{5,6}

This post hoc analysis evaluated the effect of imetelstat on proinflammatory cytokines and the impact on efficacy in patients from IMbark

Methods

- The study design for IMbark is presented in **Figure 1**. Patients were randomly assigned (1:1) to 4.4 mg/kg or 8.9 mg/kg imetelstat active dose (equivalent to 4.7 mg/kg or 9.4 mg/kg imetelstat sodium, respectively) by infusion every 3 weeks
- In this analysis, 21 cytokines were measured using Luminex® Multiplex Cytokine Assay Kits (R&D Systems, Inc.; Minneapolis, MN) in stored plasma samples collected at baseline and at weeks 12, 24, 36, and 48 of imetelstat treatment
- The correlation between percent change from baseline in cytokine levels and percent change in total symptom score (TSS) or spleen volume at each time point was evaluated using linear regression
- OS with imetelstat (any dose or by dose level) by baseline cytokine level was estimated using Kaplan-Meier; hazard ratio (HR) between cytokine levels was evaluated using Cox regression
- An unadjusted nominal *P* value from each test was reported

Figure 1. IMbark Study Design




CR, complete remission; IWG-MRT, International Working Group-Myeloproliferative Neoplasms Research and Treatment; JAKi, Janus kinase inhibitor; OS, overall survival; PR, partial remission; R, randomized; R/R, relapsed or refractory; SVR, spleen volume reduction; TSS, total symptom score.

- For R/R status, patients must have had worsening of splenomegaly-related abdominal pain at any time after the start of JAKi therapy either:
 - No reduction in spleen volume or size after 12 weeks of JAKi therapy
 - Worsening splenomegaly after starting JAKi, with documented spleen volume increase by 25% or increase in spleen size by palpation

Results


- A total of 43 out of 107 patients from IMbark who had plasma samples available for cytokine evaluation and quantifiable cytokine levels at baseline for ≥1 cytokine and ≥2 postbaseline assessments at 12 and 24 weeks were included in this analysis
- Of 21 cytokines analyzed, a consistent trend of dose-dependent reduction across time points was seen selectively with IL-8 and TNF-α levels (**Figure 2** and **Figure 3**)

Figure 2. Mean Percentage Change From Baseline in 21 Cytokines in Patients Treated With (A) 4.4 mg/kg Imetelstat or (B) 8.9 mg/kg Imetelstat Through Week 48



C, cycle; D, day; GM-CSF, granulocyte-macrophage colony-stimulating factor; IFN, interferon; IL, interleukin; ITAC, interleukin-1 receptor-associated kinase 1; MIP, macrophage inflammatory protein; TNF-α, tumor necrosis factor alpha.
Note: A visit window of ±1 cycle was applied to derive cycles 5, 9, 13, and 17 for cytokine data, corresponding to weeks 12, 24, 36, and 48 (±3 weeks).
*Only 1 patient in the 4.4-mg/kg cohort and 4 patients in the 8.9-mg/kg cohort had detectable MIP-1α results.

Figure 3. Mean Percentage Change From Baseline in (A) IL-8 and (B) TNF-α Levels Through Week 48 in Patients Treated With Imetelstat




C, cycle; D, day; IL, interleukin; TNF-α, tumor necrosis factor alpha.
Note: A visit window of ±1 cycle was applied to derive cycles 5, 9, 13, and 17 for cytokine data, corresponding to weeks 12, 24, 36, and 48 (±3 weeks).

Response Outcomes


- Reductions from baseline in IL-8 levels with imetelstat (regardless of dose) at weeks 12, 24, and 36 were significantly correlated with reductions in TSS at the corresponding time points (correlation coefficients [CC] of 0.361 [*P*=.019], 0.319 [*P*=.045], and 0.349 [*P*=.040], respectively) and with spleen volume reduction (SVR) at week 12 (CC, 0.399; *P*=.009) (**Figure 4**)
 - The correlations between changes in IL-8 and in SVR at weeks 24 and 36 were not statistically significant (**not shown**)
- Reductions from baseline in TNF-α levels with imetelstat (regardless of dose) at weeks 12, 24, 36, and 48 were significantly correlated with SVR at the corresponding time points (CCs of 0.636 [*P*<.001], 0.457 [*P*=.003], 0.439 [*P*=.011], and 0.475 [*P*=.012], respectively) (**Figure 5**)
 - No statistically significant correlations between the changes in TNF-α and in TSS were observed (**not shown**)

Figure 4. Change in IL-8 From Baseline With Imetelstat Is Correlated With Change in TSS at Weeks (A) 12, (B) 24, and (C) 36, and With (D) Change in SVR at Week 12



IL, interleukin; IR, independent review committee; SVR, spleen volume reduction; TSS, total symptom score.


Figure 5. Change in TNF-α From Baseline With Imetelstat Is Correlated With Change in SVR at Weeks (A) 12, (B) 24, (C) 36, and (D) 48



IR, independent review committee; SVR, spleen volume reduction; TNF-α, tumor necrosis factor alpha.

- Patients who achieved spleen response (≥35% SVR) at week 24 with imetelstat had mean reductions of 32% and 46%, respectively, in IL-8 and TNF-α levels at week 24; SVR nonresponders had mean reductions of 2% and 13%, respectively (**Figure 6**)

Figure 6. Greater Reduction From Baseline in (A) IL-8 and (B) TNF-α Levels in SVR Responders Than in Nonresponders at Week 24




IL, interleukin; SVR, spleen volume reduction; TNF-α, tumor necrosis factor alpha.

Survival Outcomes

- Overall, patients with lower baseline IL-8 levels (defined as ≤ median level) had longer median OS with imetelstat compared with patients with higher baseline IL-8 levels (defined as > median level; HR [95% CI], 0.44 [0.21-0.93]; *P*=.03) (**Figure 7A**)
 - Among patients with higher baseline IL-8 levels, median OS (95% CI) was longer with 8.9 mg/kg imetelstat (24.1 months [18.4-31.6]) compared with 4.4 mg/kg imetelstat (18.2 months [14.7-not estimable]; HR [95% CI], 0.45 [0.14-1.48]; *P*=.18) (**Figure 7B**)
- Similar trends in OS were observed for TNF-α, but these were not statistically significant; patients with lower baseline TNF-α levels had longer median OS compared with patients with higher baseline TNF-α levels (HR [95% CI], 0.64 [0.31-1.31]; *P*=.21) (**Figure 7C**)
 - Among patients with higher baseline TNF-α levels, median OS (95% CI) was longer with 8.9 mg/kg imetelstat (22.8 months [13.9-not estimable]) compared with 4.4 mg/kg imetelstat (19.3 months [14.7-34.7]; HR [95% CI], 0.68 [0.24-1.88]; *P*=.45) (**Figure 7D**)

Figure 7. OS by (A) Baseline IL-8 Level, (B) Imetelstat Dose and Baseline IL-8 Level, (C) Baseline TNF-α Level, and (D) Imetelstat Dose and Baseline TNF-α Level



IL, interleukin; TNF-α, tumor necrosis factor alpha.

Conclusions

- In this post hoc analysis, imetelstat treatment was associated with dose-dependent reductions from baseline in IL-8 and TNF-α levels, which corresponded with reductions in TSS or spleen volume
- In a small number of patients with higher baseline IL-8 and TNF-α, 8.9 mg/kg imetelstat showed a longer survival benefit compared with 4.4 mg/kg imetelstat, despite the known association of elevated IL-8 and TNF-α with worse prognosis
- This hypothesis-generating, exploratory analysis identifies specific inflammatory cytokines reduced with imetelstat treatment and associated with reduction in spleen volume and symptoms that, together with previously presented data, suggest potential disease-modifying activity of imetelstat in MF
- Further confirmatory studies of imetelstat in MF are in progress

References

- Wang Z, et al. *Front Oncol*. 2024;14:1369346.
- Zahn SA, et al. *Hematology*. 2016;2016:1660-671.
- Teffari A, et al. *J Clin Oncol*. 2013;29(10):1356-1363.
- Solomon P, et al. *BMC Cancer*. 2018;18(1):730.
- Mascarenhas J, et al. *J Clin Oncol*. 2021;39(26):2881-2892.
- Kuykendall AT, et al. *Ann Hematol*. 2022;101(1):139-146.

Acknowledgments

- The authors thank all the patients and caregivers for their participation in this study and acknowledge the collaboration and commitment of all investigators and their research support staff
- This study was funded by Geron Corporation
- All authors contributed to and approved the presentation, writing and editorial support were provided by Jeremy J. Henriques, PhD, CMPP, of The Lockwood Group (Stamford, CT, USA), funded by Geron Corporation