

Presented at the
Society of Hematologic Oncology
(SOHO) Annual Meeting
Houston, TX, USA • September 9-12, 2026
Originally presented at EHA 2026

Haris Ali,¹ Tania Jain,² Aaron Gerdts,³ Vikas Gupta,⁴ Claire Harrison,⁵ Joan How,⁶ David M. Ross,⁷ Lucia Masarova,⁸ Bethan Psaila,⁹ Abdulraheem Yacoub,¹⁰ Terrence Bradley,¹¹ Caroline McNamara,¹² Andrew Perkins,¹³ William Shomali,¹⁴ Makoto Yoshimitsu,¹⁵ Junichiro Yuda,¹⁶ Haifa Kathrin Al-Ali,¹⁷ Lynette Chee,¹⁸ Jean-Jacques Kiladjian,¹⁹ Ruben Mesa,²⁰ Francesco Passamonti,²¹ Raajit Rampal,²² Tim Somerville,²³ Alessandro Maria Vannucchi,²⁴ Hiroki Yamaguchi,²⁵ Steffen Koschmieder,²⁶ Jason Gotlib,¹⁴ Jyoti Nangalia,²⁷ Chenwei Tian,²⁸ Betty Lamothe,²⁸ Erin Crowgey,²⁸ Tatiana Zinger,²⁸ Evan Braunstein,²⁸ John Mascarenhas²⁹

¹City of Hope Medical Center, Duarte, CA, USA; ²Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University, Baltimore, MD, USA; ³Cleveland Clinic, Cleveland, OH, USA; ⁴Princess Margaret Cancer Centre, Toronto, ON, Canada; ⁵Guy's and St Thomas' NHS Foundation Trust, London, UK; ⁶Brigham and Women's Hospital and Dana-Farber Cancer Institute, Boston, MA, USA; ⁷Royal Adelaide Hospital, Adelaide, SA, Australia; ⁸MD Anderson Cancer Center, Houston, TX, USA; ⁹University of Oxford, Oxford, UK; ¹⁰The University of Kansas Cancer Center, Kansas City, KS, USA; ¹¹University of Miami Sylvester Comprehensive Cancer Center, Division of Hematology, Miami, FL, USA; ¹²Royal Brisbane and Women's Hospital, Brisbane, QLD, Australia; ¹³The Alfred Hospital, Melbourne, VIC, Australia; ¹⁴Stanford Cancer Institute/Stanford University School of Medicine, Stanford, CA, USA; ¹⁵Kagoshima University Hospital, Kagoshima, Japan; ¹⁶Department of Hematology, National Cancer Center Hospital East, Chiba, Japan; ¹⁷University Medicine Halle, Saale, Germany; ¹⁸Peter MacCallum Cancer Centre and The Royal Melbourne Hospital, Melbourne, VIC, Australia; ¹⁹Saint-Louis Hospital, Paris Cité University, INSERM, Paris, France; ²⁰Wake Forest University Baptist Medical Center, Winston-Salem, NC, USA; ²¹Fondazione IRCCS Ca' Ganda Ospedale Maggiore, Milan, Italy; ²²Memorial Sloan Kettering Cancer Center, New York, NY, USA; ²³Leukaemia Biology Laboratory, Cancer Research UK Manchester Institute, The University of Manchester, Manchester, UK; ²⁴Azienda Ospedaliero-Universitaria Careggi, Florence, Italy; ²⁵Nippon Medical School Hospital, Tokyo, Japan; ²⁶RWTH Aachen University, Faculty of Medicine, and Center for Integrated Oncology (CIO-ABCD), Aachen, Germany; ²⁷The Sanger Institute, Cambridge, UK; ²⁸Incyte Corporation, Wilmington, DE, USA; ²⁹Icahn School of Medicine at Mount Sinai, New York, NY, USA

Introduction


Calreticulin Mutations Are Frequent in Essential Thrombocythemia With No Mutant-Specific Treatment Available

- Essential thrombocythemia (ET) is characterized by megakaryocyte hyperplasia and clustering, thrombocytosis, and an increased risk of thrombosis, hemorrhage, and transformation to myelofibrosis (MF)^{1,2}
- Mutations of calreticulin (mutCALR) in exon 9 are found in 25-30% of patients with ET³⁻⁵
- ET with mutCALR is associated with diagnosis at a younger age and higher risk of transformation to MF compared with JAK2V617F^{6,7}
- Current treatments are broadly myelosuppressive, not mutant targeted, and have limited efficacy in reducing mutCALR allele frequency^{8,9}

INCA033989 Is a mutCALR-Targeted Therapy for Patients With ET and MF

- INCA033989 has a unique mechanism of action (Figure 1)
 - INCA033989 is a novel, first-in-class, fully human, Fc-silenced, immunoglobulin G1 monoclonal antibody that selectively targets mutCALR with high affinity, when in complex with thrombopoietin receptor, to inhibit oncogenic signaling and proliferation of cells¹⁰


Figure 1. INCA033989 Mechanism of Action



Methods

- INCA033989-101** (NCT05936359; outside the US) and **INCA033989-102** (NCT06034002; US only) are phase 1, first-in-human, multicenter, open-label studies evaluating INCA033989 in patients harboring a CALR exon-9 mutation with high-risk ET or MF (as monotherapy or in combination with ruxolitinib)
- INCA033989 was administered intravenously every 2 weeks (range 24-2500 mg), and alternative frequencies were tested every 4 weeks, other; Figure 2)

Figure 2. Study Design



ELN, European LeukemiaNet; ET, essential thrombocythemia; JAK1, Janus kinase inhibitor; MF, myelofibrosis; MPN-SAF, Myeloproliferative Neoplasms Symptom Assessment Form; mutCALR, mutations of calreticulin; TSS, total symptom score; WHO, World Health Organization.

Results

Table 1. Demographics and Disease Characteristics	
Variable	Total (N=114)
Age, years, median (range)	62 (23, 89)
≥65, n (%)	48 (42.1)
Female, n (%)	69 (60.5)
Time from initial diagnosis, years, median (range)	6.9 (0.3, 27.9)
CALR exon 9 mutation type, n (%)	
Type 1	50 (43.9)
Non-Type 1	64 (56.1)
Median CALR VAF,* % (range)	34 (10, 75)
Median platelets (range), G/L	913 (444, 3152)
Receiving anticoagulant therapy at baseline, n (%)	17 (14.9)
Receiving aspirin at baseline, n (%)	69 (60.5)
Prior cytoreductive therapy†, n (%)	
Hydroxycarbamide	100 (87.7)
Anagrelide	36 (31.6)
Interferons	35 (30.7)
Other‡	12 (10.5)

Data cutoff: March 6, 2026.
*Measured centrally in peripheral blood by NGS (n=107). †Categories not mutually exclusive. ‡bomedsstat (n=2), pelabresb (n=1), busulfan (n=2), ruxolitinib (n=9), CALR, calreticulin; NGS, next generation sequencing; VAF, variant allele frequency.

INCA033989 Monotherapy Is Well Tolerated

- 95% (108/114) of patients remain on treatment; only 5% (6/114) of patients discontinued treatment
- The median (range) duration of INCA033989 exposure was 8.1 months (0.59, 27.0)
- Grade ≥3 cytopenia treatment-emergent adverse events (TEAEs) occurred in 6% (7/114) of patients; no grade ≥3 thrombocytopenia TEAEs were observed (Table 2)
- Lipase or amylase elevations were transient, with no associated clinical sequelae

Table 2. Safety Outcomes	
Overall TEAE Summary	Most Common TEAEs (≥10% of Patients)
TEAE, n (%)	TEAE, n (%)
Any TEAE	Fatigue
Treatment-related	Headache
Grade ≥3*	Lipase increased
Serious†	Constipation
Fatal	Diarrhea
Discontinuation due to TEAEs	Anemia
Dose reduction due to TEAEs	Nausea
Infusion interruption due to TEAEs	URT†
Dose delay due to TEAEs	Dizziness
Dose-limiting toxicity	Arthralgia
	Pruritus
	Amylase increased

*One grade 4 TEAE was observed (transient neutropenia related to concomitant hydroxyurea). †The following serious adverse events were considered unrelated to INCA033989: acute sinusitis (n=1; 750 mg), basal cell carcinoma (n=1; 2500 mg Q2W), diverticulitis (n=1; 400 mg), polymyalgia rheumatica (n=1; 1500 mg), proctitis (n=1; 1500 mg), urinary tract infection (n=1; 1500 mg). One event, visceral venous thrombosis (n=1; 24 mg) followed by melena (after anticoagulant initiation) and treatment discontinuation, was related to INCA033989. ‡Adverse event (visceral venous thrombosis, n=1), lack of efficacy (n=2), pregnancy (n=1), and patient withdrawal (n=2). Q2W, every 2 weeks. TEAE, treatment-emergent adverse event; URT†, upper respiratory tract infection.

Figure 3. Platelet Count Response by Dose and mutCALR Type (n=63)

